

AD-A151 070

CONVOLUTION OF THE IFRA (INCREASING FAILURE RATE  
AVERAGE) SCALED-MINS CLA. (U) PITTSBURGH UNIV PA DEPT  
OF MATHEMATICS AND STATISTICS E EL-NEWELHI ET AL.

1/1

UNCLASSIFIED


JAN 85 AFOSR-TR-85-0109 N00014-84-K-0084

F/G 12/1

NL

END

FIGURE 10



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

Contract no. N00014-84-K-0084 verified  
per telecom to Office of Naval Research.

3-15-85 slb

**UNCLASSIFIED**

SECURITY CLASSIFICATION OF TITLE

1b. REPORT SECURITY CLASSIFICATION  
**UNCLASSIFIED**

2a. SECURITY CLASSIFICATION

2b. DECLASSIFICATION/DOWNGRADING

4. PERFORMING ORGANIZATION  
U of I 85-1

6a. NAME OF PERFORMING ORGANIZATION  
University of Illinois

6c. ADDRESS (City, State and ZIP)  
Department of Mathematics  
Computer Science, Chicago

8a. NAME OF FUNDING/SPONSORING ORGANIZATION  
AFOSR

8c. ADDRESS (City, State and ZIP)

Bolling AFB DC 2033

11. TITLE (Include Security Classification)  
CONVOLUTION OF THE I

12. PERSONAL AUTHOR(S)  
Emad El-Newehi and

13a. TYPE OF REPORT  
Technical

16. SUPPLEMENTARY NOTATION

17. COSATI CODES	
FIELD	GROUP

19. ABSTRACT (Continue on reverse)

The class of non  
is IFRA for all

20. DISTRIBUTION/AVAILABILITY STATEMENT  
UNCLASSIFIED/UNLIMITED ☒

22a. NAME OF RESPONSIBLE PERSON  
MAJ Brian W. Woodruff

DD FORM 1473, 83 APR

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

## REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION <b>UNCLASSIFIED</b>			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited.		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) U of I 85-1			5. MONITORING ORGANIZATION REPORT NUMBER(S) <b>AFOSR-TR- 85 - 0109</b>		
6a. NAME OF PERFORMING ORGANIZATION Univeristy of Illinois		6b. OFFICE SYMBOL (If applicable)		7a. NAME OF MONITORING ORGANIZATION Air Force Office of Scientific Research	
6c. ADDRESS (City, State and ZIP Code) Department of Mathematics, Statistics, and Computer Science, Chicago IL 60607			7b. ADDRESS (City, State and ZIP Code) Directorate of Mathematical & Information Sciences, Bolling AFB DC 20332-6448		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION AFOSR		8b. OFFICE SYMBOL (If applicable) NM		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER AFOSR-80-0170	
8c. ADDRESS (City, State and ZIP Code) Bolling AFB DC 20332-6448			10. SOURCE OF FUNDING NOS.		
			PROGRAM ELEMENT NO. 61102F	PROJECT NO. 2304	TASK NO. A5
11. TITLE (Include Security Classification) CONVOLUTION OF THE IFRA SCALED-MINS CLASS					
12. PERSONAL AUTHOR(S) Emad El-Newehi and Thomas H. Savits					
13a. TYPE OF REPORT Technical		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Yr., Mo., Day) JAN 85	
15. PAGE COUNT 6					
16. SUPPLEMENTARY NOTATION					
17. COSAFI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB. GR.	Increasing failure rate average; characterizations; convolution, Equations. < _____		
19. ABSTRACT (Continue on reverse if necessary and identify by block number)					
<p>The class of nonnegative random vectors <math>I = (T_1, \dots, T_n)</math> for which <math>\min_{1 \leq i \leq n} a_i T_i</math> is IFRA for all <math>0 &lt; a_i \leq \infty</math>, <math>i = 1, \dots, n</math>, is closed under convolution.</p>					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS <input type="checkbox"/>			21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED		
22a. NAME OF RESPONSIBLE INDIVIDUAL MAJ Brian W. Woodruff			22b. TELEPHONE NUMBER (Include Area Code) (202) 767- 5027		22c. OFFICE SYMBOL NM

CONVOLUTION OF THE IFRA SCALED-MINS CLASS

by

Emad El-Neweih<sup>1,3</sup>

and

Thomas H. Savits<sup>2,4</sup>

DTIC  
ELECTE  
MAR 6 1985  
S B D

University of Illinois at Chicago

Department of Mathematics, Statistics, and Computer Science

Technical Report No. 85-1

Statistical Laboratory

Grant no. AFOSR-84-0113 verified by January, 1985  
Work Unit Search.

Performing Organization - Univ. of  
Pittsburgh. slb 3-15-85

Contract no. N00014-84-K-0084 verified  
per telecom to Office of Naval Research.  
Contractor - Univ. of Pittsburgh.  
3-15-85 slb

<sup>1</sup>Department of Mathematics, Statistics and Computer Science, University of  
Illinois at Chicago, Chicago, IL

<sup>2</sup>Department of Mathematics and Statistics, University of Pittsburgh, Pittsburgh,  
PA 15260.

<sup>3</sup>Supported by Grant No. AFOSR-80-0170 and Grant No. AFOSR-84-0113.

<sup>4</sup>Supported by Grant No. AFOSR-84-0113 and ONR Contract N00014-84-K-0084.

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)

NOTICE OF TRANSMITTAL TO DTIC

This technical report has been reviewed and is  
approved for public release IAW AFR 190-12.

Distribution is unlimited.

MATTHEW J. KIRPER

Chief, Technical Information Division

# Convolution of the IFRA Scaled-mins Class

By

Emad El-Newehi

and

Thomas H. Savits

## Abstract

The class of nonnegative random vectors  $\underline{T} = (T_1, \dots, T_n)$  for which  $\min_{1 \leq i \leq n} a_i T_i$  is IFRA for all  $0 < a_i \leq \infty$ ,  $i = 1, \dots, n$ , is closed under convolution.

**Key Words and Phrases:** Increasing failure rate average, characterizations, convolution.

**AMS 1980 subject classifications:** Primary 62H05, Secondary 62N05.

Accession For	
NTIS	<input checked="" type="checkbox"/>
DTIC Tab	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution	
Availability Codes	
Avail and/or	
Dist	Special
A-1	



## 1. Introduction and statement of main result.

*5/2/84* In recent years various multivariate extensions of the univariate classes of life distributions that are important in reliability theory have been proposed. A survey of many of these classes may be found in Block and Savits (1981). In this paper we focus on one particular extension of the IFRA (increasing failure rate average) class due to Esary and Marshall (1979). *key words include: 4/2* nonnegative random vector  $\underline{T} = (T_1, \dots, T_n)$  is said to satisfy condition (F) if  $\min_{1 \leq i \leq n} a_i T_i$  is IFRA for all choices  $0 < a_i < \infty$ ,  $i = 1, \dots, n$ . (Recall that a nonnegative random variable  $T$  is IFRA if  $\bar{F}(\alpha t) \geq \bar{F}(t)$  for all  $t \geq 0$ ,  $0 < \alpha < 1$ , where  $\bar{F}(t) = P(T > t)$  is the survival probability). Here we interpret  $\infty \cdot 0 = \infty$ .

Although Esary and Marshall (1979) considered some closure properties of this class, they did not study closure with respect to the operation of convolution. Recently, El-Newehi (1984) showed that the class is closed under convolution provided one of the two vectors has independent components; however, the general problem was not resolved. The purpose of this paper is to prove the general result as stated below.

Let  $I$  denote the class of all nonnegative random vectors  $\underline{T} = (T_1, \dots, T_n)$  satisfying condition (F).

Theorem 1.1. The class  $I$  is closed under convolution.

The proof of this result is contained in section 2. In section 3 we consider a further characterization of the class  $I$ . As usual  $R_+^n$  denotes the nonnegative upper orthant.

## 2. Proof of the main result.

We shall make use of the following characterization of the class  $I$  due to El-Newehi (1984).

**Theorem 2.1** (El-Newehi). A nonnegative random vector  $\underline{T} = (T_1, \dots, T_n)$  belongs to  $I$  if and only if

$$E\left[\prod_{i=1}^n h_i(T_i)\right] \leq E^{1/\alpha}\left[\prod_{i=1}^n h_i^\alpha(T_i/\alpha)\right] \quad (2.1)$$

for all  $0 < \alpha < 1$  and all nonnegative nondecreasing functions  $h_i$  defined on  $[0, \infty)$ ,  $i = 1, \dots, n$ .

Thus to show that  $I$  is closed under convolution, we need only show that if  $\underline{S} = (S_1, \dots, S_n)$  and  $\underline{T} = (T_1, \dots, T_n)$  are independent vectors in  $I$ , then

$$E\left[\prod_{i=1}^n h_i(S_i + T_i)\right] \leq E^{1/\alpha}\left[\prod_{i=1}^n h_i^\alpha((S_i + T_i)/\alpha)\right]$$

for all  $0 < \alpha < 1$  and all nonnegative nondecreasing  $h_i$  on  $[0, \infty)$ ,  $i = 1, \dots, n$ .

As usual we may assume without loss of generality that each  $h_i$  is continuous and bounded (see, e.g., Block and Savits (1980)).

First we prove a lemma.

**Lemma 2.2.** Let  $H(\underline{s}, \underline{t})$  be bounded, nonnegative and continuous on  $\mathbb{R}_+^n \times \mathbb{R}_+^n$ . Let  $\mu$  and  $\nu$  be two finite measures on  $\mathbb{R}_+^n$ . For  $0 < \alpha < 1$ , define  $\|H(\cdot, \underline{t})\|_\alpha = \left\{ \int H^\alpha(\underline{s}, \underline{t}) d\nu(\underline{s}) \right\}^{1/\alpha}$ . Then

$$\begin{aligned} \int \|H(\cdot, \underline{t})\|_\alpha d\mu(\underline{t}) &\leq \left\{ \int \left[ \int H(\underline{s}, \underline{t}) d\mu(\underline{t}) \right]^\alpha d\nu(\underline{s}) \right\}^{1/\alpha} \\ &= \left\| \int H(\cdot, \underline{t}) d\mu(\underline{t}) \right\|_\alpha. \end{aligned} \quad (2.2)$$

**Proof.** If  $m > 0$  and  $\underline{i} = (i_1, \dots, i_n)$ , let  $A_{\underline{i}}^m = \left[ \frac{i_1-1}{2^m}, \frac{i_1}{2^m} \right) \times \dots \times \left[ \frac{i_n-1}{2^m}, \frac{i_n}{2^m} \right)$  for  $1 \leq i_j \leq m \cdot 2^m$ ,  $j = 1, \dots, n$ ,  $m = 1, 2, \dots$ . Set  $H_m(\underline{s}, \underline{t}) = H(\underline{s}, 2^{-m}\underline{i})$  for  $\underline{t} \in A_{\underline{i}}^m$  and zero otherwise. Since  $H$  is bounded and continuous,  $H_m(\underline{s}, \underline{t}) \rightarrow H(\underline{s}, \underline{t})$  boundedly as  $m \rightarrow \infty$ . Hence  $\|H_m(\cdot, \underline{t})\|_\alpha \rightarrow \|H(\cdot, \underline{t})\|_\alpha$  boundedly and



$\int \|H_m(\cdot, \underline{t})\|_\alpha d\mu(\underline{t}) \rightarrow \int \|H(\cdot, \underline{t})\|_\alpha d\mu(\underline{t})$  as  $m \rightarrow \infty$  by the bounded convergence theorem. But

$$\begin{aligned} \int \|H_m(\cdot, \underline{t})\|_\alpha d\mu(\underline{t}) &= \sum_{\underline{1}} \|H_m(\cdot, 2^{-m}\underline{1})\|_\alpha \mu(A_{\underline{1}}^m) \\ &= \sum_{\underline{1}} \|\mu(A_{\underline{1}}^m) H_m(\cdot, 2^{-m}\underline{1})\|_\alpha \\ &\leq \left\| \sum_{\underline{1}} \mu(A_{\underline{1}}^m) H_m(\cdot, 2^{-m}\underline{1}) \right\|_\alpha \\ &= \left\| \int H_m(\cdot, \underline{t}) d\mu(\underline{t}) \right\|_\alpha. \end{aligned}$$

The inequality follows from Minkowski's inequality for  $0 < \alpha < 1$ . The desired result is obtained by passing to the limit as  $m \rightarrow \infty$ .

(2.3) Remark. The above lemma remains valid if we weaken the continuity assumption on  $H$ . It suffices that  $H(\underline{s}, \underline{t})$  be measurable and right-continuous in  $\underline{t}$  for each fixed  $\underline{s}$ . We can also replace right-continuity with left-continuity if we redefine  $H_m(\underline{s}, \underline{t})$  as  $H(\underline{s}, 2^{-m}(\underline{1}-\underline{1}))$  on  $A_{\underline{1}}^m$  where  $\underline{1} = (1, \dots, 1)$ .

We are now ready to prove the main result. Let  $\underline{S} = (S_1, \dots, S_n)$  and  $\underline{T} = (T_1, \dots, T_n)$  be independent vectors in  $I$  with corresponding distribution functions  $F$  and  $G$  respectively. Fix  $0 < \alpha < 1$  and let  $h_i$  be nonnegative, non-decreasing, continuous bounded functions on  $[0, \infty)$ . Then

$$\begin{aligned} E\left[\prod_{i=1}^n h_i(S_i + T_i)\right] &= \iint \prod_{i=1}^n h_i(s_i + t_i) dF(\underline{s}) dG(\underline{t}) \\ &\leq \int \left[ \int \prod_{i=1}^n h_i\left(\frac{s_i}{\alpha} + t_i\right) dF(\underline{s}) \right]^{1/\alpha} dG(\underline{t}) \quad (\text{since } \underline{S} \in I) \\ &\leq \left\{ \int \left[ \int \prod_{i=1}^n h_i\left(\frac{s_i}{\alpha} + t_i\right) dG(\underline{t}) \right]^\alpha dF(\underline{s}) \right\}^{1/\alpha} \quad (\text{By Lemma 2.2}). \end{aligned}$$

$$\begin{aligned} &\leq \left\{ \left[ \left( \prod_{i=1}^n h_i^\alpha \left( \frac{s_i}{\alpha} + \frac{t_i}{\alpha} \right) dG(\underline{t}) \right)^{1/\alpha} dF(\underline{s}) \right]^{1/\alpha} \quad (\text{since } \underline{T} \in I) \\ &= E^{1/\alpha} \left[ \prod_{i=1}^n h_i^\alpha \left( \frac{S_i + T_i}{\alpha} \right) \right]. \end{aligned}$$

(2.4) Remark. Suppose  $H$  is any class of nonnegative functions and we define  $I$  to be  $H$ -IFRA if  $E[h(\underline{T})] \leq E^{1/\alpha} [h^\alpha(\underline{T}/\alpha)]$  for all  $h \in H$ ,  $0 < \alpha < 1$ . Then this same argument shows that such a class is closed under convolution provided whenever  $h \in H$ , it follows that  $h(\underline{s} + \underline{t})$  belongs to  $H$  for fixed  $\underline{s}$  and for fixed  $\underline{t}$ .

### 3. Another characterization of $I$

As was mentioned in Section 2, El-Newehi characterized the class  $I$  by the requirement that

$$E[H(\underline{T})] \leq E^{1/\alpha} [H^\alpha(\underline{T}/\alpha)] \quad (3.1)$$

for all  $0 < \alpha < 1$  and  $H(\underline{t})$  of the form  $\prod_{i=1}^n h_i(t_i)$ , where each  $h_i$  is nonnegative and nondecreasing on  $[0, \infty)$ . With the help of Lemma 2.2 we can extend the inequality (3.1) to a larger class.

Let  $H$  denote the class of all nonnegative distribution functions on  $R_+^n$ ; i.e.,  $H \in H$  if and only if there exists a measure  $\mu$  on  $R_+^n$  such that  $H(\underline{t}) = \mu([0, \underline{t}])$ . We denote this unique measure  $\mu$  by  $dH$ .

Theorem 3.2.  $\underline{T} \in I$  if and only if

$$E[H(\underline{T})] \leq E^{1/\alpha} [H^\alpha(\underline{T}/\alpha)]$$

for all  $0 < \alpha < 1$  and all  $H \in H$ .

Proof. The sufficiency is clear since  $h(\underline{t}) = \prod_{i=1}^n h_i(t_i) \in H$  whenever each  $h_i$  is nonnegative, nondecreasing and right-continuous. Now suppose  $\underline{T} \in I$  and  $H \in H$ . Let  $F$  be the distribution of  $\underline{T}$ . Then

$$\begin{aligned}
E[H(\underline{T})] &= \int H(\underline{t}) dF(\underline{t}) = \iint I_{[0, \underline{t}]}(\underline{s}) dH(\underline{s}) dF(\underline{t}) \\
&= \int \left[ \int I_{[\underline{s}, \infty)}(\underline{t}) dF(\underline{t}) \right] dH(\underline{s}) \\
&\leq \int \left[ \int I_{[\underline{s}, \infty)}(\underline{t}/\alpha) dF(\underline{t}) \right]^{1/\alpha} dH(\underline{s}) && (\text{since } \underline{T} \in I) \\
&\leq \left\{ \int \left[ \int I_{[0, \underline{t}/\alpha]}(\underline{s}) dH(\underline{s}) \right]^\alpha dF(\underline{t}) \right\}^{1/\alpha} && (\text{Remark (2.3)}) \\
&= E^{1/\alpha} [H^\alpha(\underline{T}/\alpha)].
\end{aligned}$$

(3.3) Remark. The characterization of the NBU class considered by El-Newehi (1984) also extends to this class  $H$ .

- [1] Block, H.W. and Savits, T.H. (1981). Multivariate classes in reliability theory. Math. of O.R., 6, 453-461.
- [2] Block, H.W. and Savits, T.H. (1980). Multivariate increasing failure rate average distributions. Ann. Prob., 8, 793-801.
- [3] El-Newehi, E. (1984). Characterizations and closure under convolution of two classes of multivariate life distributions. Statistics and Probability Letters. To appear.
- [4] Esary, J.D. and Marshall, A.W. (1979). Multivariate distributions with increasing hazard rate averages. Ann. Prob., 7, 359-370.

**END**

**FILMED**

**4-85**

**DTIC**